



Character Formulas from Lusztig Varieties and Affine Springer Fibers

Minh-Tâm Quang Trinh

Yale University

This talk is about...

- 1 Braids
- 2 Lusztig Varieties
- 3 Springer Fibers
- 4 Affine Springer Fibers

1 Braids The braid group $Br_n =$


$$\left\langle \sigma_1, \dots, \sigma_{n-1}, \left| \begin{array}{l} \sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1}, \\ \sigma_i \sigma_j = \sigma_j \sigma_i \text{ when } |i - j| > 1 \end{array} \right. \right\rangle$$

appears in [knot theory](#) and [representation theory](#).

$$\begin{array}{c} \sigma_i \\ \left[\dots \right] \sum_{i \quad i+1} \left[\dots \right] \end{array}$$

A *link* is a collection of circles (tamely) embedded in \mathbf{R}^3 . Knot theory is about isotopy invariants of links.

(Alexander) Every link is the *closure* of some braid.



Let $G = \mathrm{GL}_n$ and B its upper-triangular subgroup.

$$V_n(q) = \{\text{functions } G(\mathbf{F}_q)/B(\mathbf{F}_q) \rightarrow \mathbf{C}\},$$

$$H_n(q) = \mathrm{End}_{G(\mathbf{F}_q)}(V_n(q)).$$

$$\text{(Iwahori)} \quad H_n(q) \simeq \frac{\mathbf{C}Br_n}{\langle \sigma_i^2 - (q-1)\sigma_i - q \rangle}.$$

To explain, recall Bruhat: $G = \coprod_{w \in S_n} B\dot{w}B$.

Then $\mathbf{C}Br_n \curvearrowright V_n(q)$ via

$$\sigma_i \cdot \mathbf{1}_{xB(\mathbf{F}_q)} = \sum_{xB \xrightarrow{i} yB} \mathbf{1}_{yB(\mathbf{F}_q)},$$

where $xB \xrightarrow{i} yB$ means $Bx^{-1}yB = B\dot{w}_{(i,i+1)}B$.

Motivates a Hecke algebra $H_n(\mathfrak{q})$ over $\mathbf{C}[\mathfrak{q}^{\pm 1}]$.

Oceanu used functions $Br_n \rightarrow H_n(\mathfrak{q}) \rightarrow \mathbf{C}(\mathfrak{q}^{\frac{1}{2}})[a^{\pm 1}]$ to construct a link invariant

$$\text{HOMFLYPT} : \{\text{links in } \mathbf{R}^3\}/\text{isotopy} \rightarrow \mathbf{C}(\mathfrak{q}^{\frac{1}{2}})[a^{\pm 1}].$$

Jones computed it for torus knots. Remarkably, the values encode \mathfrak{q} -Catalan (and \mathfrak{q} -Kirkman) numbers.

On the other hand, Iwahori suggests that HOMFLYPT is related to the geometry of G/B .

We'll discuss a Springer-theoretic function of β that refines the HOMFLYPT invariant of its closure $\hat{\beta}$.

2 Lusztig Varieties Suppose that β is *positive*:

$$\beta = \sigma_{i_1} \cdots \sigma_{i_\ell}.$$

(Deligne) The variety $O(\beta) =$

$$\left\{ (g_0 B, g_1 B, \dots, g_\ell B) \mid g_{j-1} B \xrightarrow{i_j} g_j B \text{ for all } j \right\}$$

only depends on β , up to isomorphisms that keep $g_0 B$ and $g_\ell B$ fixed.

For any positive β, β' , we have

$$O(\beta\beta') \simeq O(\beta) \times_{G/B} O(\beta'),$$

where $\times_{G/B}$ means the variety of pairs $(\vec{g}B, \vec{g}'B)$ such that $g_\ell B = g'_0 B$.

A literal geometric representation of positive braids.

For any $x \in G(\mathbf{F}_q)$, form the *braid Lusztig variety*

$$\mathcal{B}(\beta)_x = \{\vec{g}B \in O(\beta) \mid g_\ell B = xg_0 B\}.$$

(Shende–Treumann–Zaslow) Up to a monomial in $q^{\frac{1}{2}}$,

$$\frac{|\mathcal{B}(\beta)_1(\mathbf{F}_q)|}{|\mathrm{PGL}_n(\mathbf{F}_q)|}$$

is the “highest” a -degree of $\mathrm{HOMFLYPT}(\hat{\beta})$ at $\mathfrak{q} \rightarrow q$.

Example Let $n = 2$ and $\beta = \sigma_1^3 \in Br_2$.

Then $\mathrm{HOMFLYPT}(\hat{\beta}) = a^2(\mathfrak{q} + \mathfrak{q}^{-1}) - a^4$.

$$O(\beta) \simeq \{\vec{g} \in (\mathbf{P}^1)^4 \mid g_0 \neq g_1 \neq g_2 \neq g_3\},$$

$$\mathcal{B}(\beta)_1 \simeq \{\vec{g} \in (\mathbf{P}^1)^3 \mid g_1, g_2, g_3 \text{ pairwise distinct}\}.$$

PGL_2 acts simply transitively on the latter.

3 Springer Fibers How to access other a -degrees?

Let $\mathcal{U} \subseteq G$ be the unipotent variety. Observe that

$$\mathcal{B}_x := \mathcal{B}(1)_x = \{gB \mid gB = xgB\}$$

is the usual Springer fiber over x , whose cohomology defines a character of S_n :

$$\Psi_x(w) := \sum_i \mathfrak{q}^{i \operatorname{tr}(w \mid H^{2i}(\mathcal{B}_x))}.$$

Thm 1 (T) Let

$$\Psi_\beta(w) = \sum_{u \in \mathcal{U}(\mathbf{F}_q)} \frac{|\mathcal{B}(\beta)_u(\mathbf{F}_q)|}{|\operatorname{PGL}_n(\mathbf{F}_q)|} \Psi_u(w)|_{\mathfrak{q} \rightarrow q}.$$

Recall $\operatorname{Irr}(S_n) = \{\chi_\lambda \mid \lambda \vdash n\}$.

Then $(\chi_{(n-k, 1, \dots, 1)}, \Psi_\beta)_{S_n}$ sees the k th a -degree.

Think of $\beta \mapsto \Psi_\beta$ as a function

$$Br_n \rightarrow H_n(q) \rightarrow \{\text{characters of } S_n\}.$$

Closely related to work of Lusztig–Abreu–Nigro.

Example Again, let $n = 2$ and $\beta = \sigma_1^3 \in Br_2$.

Recall $\operatorname{HOMFLYPT}(\hat{\beta}) = a^2(\mathfrak{q} + \mathfrak{q}^{-1}) - a^4$.

$$\Psi_u = \begin{cases} 1 + \mathfrak{q} \operatorname{sgn} & u = 1, \\ 1 & u \neq 1. \end{cases}$$

$$\Psi_\beta = q^2 + 1 + q \operatorname{sgn}.$$

Thm 2 (T) The cohomology of $\mathcal{U}(\beta) \times_{\mathcal{U}} \mathcal{U}(1)$, where

$$\mathcal{U}(\beta) = \{(u, \vec{g}B) \mid u \in \mathcal{U} \text{ and } \vec{g}B \in \mathcal{B}(\beta)_u\},$$

encodes finer invariants of $\hat{\beta}$.

The *full twist* $\pi = (\sigma_1 \cdots \sigma_{n-1})^n$:



Thm 3 (T) Suppose $\beta^m = \pi^d$ for some $d, m > 0$.

Then up to a monomial, $\Psi_\beta(w)$ is the $\mathfrak{q} \rightarrow q$ limit of

$$\frac{\text{sgn}(w)}{\det(1 - \mathfrak{q}w \mid \mathfrak{h})} \sum_{\lambda \vdash n} \mathfrak{q}^{c(\lambda)d/m} D_\lambda(e^{2\pi id/m}) \chi_\lambda(w)$$

where:

- \mathfrak{h} is the *reflection representation*.
- $c(\lambda)$ is the sum of *contents* of λ .
- $D_\lambda(t) = K_{\lambda, (1^n)}(t)$ is the *fake degree* of λ .

Subsumes Jones's HOMFLYPT formula for torus knots.

Thm 3 generalizes to any reductive G , once we replace:

- S_n with the Weyl group W .
- $c(\lambda)$ with $c(\chi) = \sum_{\text{refl. } t} \frac{\chi(t)}{\chi(1)}$.
- fake degrees D_λ with *generic degrees* D_χ .

If $\gcd(d, m) = 1$ and m is the Coxeter number of W , then the formula simplifies:

$$\left. (\text{monomial}) \cdot \frac{\det(1 - \mathfrak{q}^d w \mid \mathfrak{h})}{\det(1 - \mathfrak{q}w \mid \mathfrak{h})} \right\} =: \Pi_{\mathfrak{q}}^{(d)}.$$

$\Pi_{\mathfrak{q}}^{(d)}$ is the character of a *rational parking space*.

$(\text{triv}, \Pi_{\mathfrak{q}}^{(d)})_W$ is a *rational q -Catalan number*.

Example If $W = S_n$, then $(\text{triv}, \Pi_{\mathfrak{q}}^{(d)})_W = \frac{[n+d-1]!}{[n]![d]!}$.

4 Affine Springer Fibers

Rational parking spaces appear in a loop or *affine* analogue of Springer theory.

finite Springer	affine Springer
G	$G((z))$
G/B	$G((z))/I$
W	$\widetilde{W} = W \ltimes X^\vee$

Above:

- $G((z))$ is the loop group $G((z))(R) := G(R((z)))$.
- I is the preimage of B in $G[[z]]$.
- X^\vee is the cocharacter lattice of $T \subseteq B$.

Dream Braid Lusztig varieties know about affine Springer representations.

We now study Springer fibers over the Lie algebras, not the groups, and over \mathbf{C} , not \mathbf{F}_q .

$$x : \mathcal{B}_x = \{gB \in G/B \mid g^{-1}xg \in \mathfrak{b}\},$$

$$\gamma = \gamma(z) : \mathcal{B}_\gamma^{\text{aff}} = \{gI \in G((z))/I \mid g^{-1}\gamma g \in \mathfrak{J}\}.$$

The table hides key differences:

In the **finite** case, \mathcal{B}_x is most **interesting** for x nilpotent.

In the **affine** case, $\mathcal{B}_\gamma^{\text{aff}}$ is *terribly infinite* for $\gamma = \gamma(z)$ nilpotent, but **interesting** for $\gamma(z)$ regular semisimple.

Example If $G = \text{SL}_2$ and $\gamma(z) = \begin{pmatrix} 0 & 1 \\ z^3 & 0 \end{pmatrix}$, then

$$\mathcal{B}_\gamma^{\text{aff}} \simeq \mathbf{P}^1 \sqcup_{\text{pt}} \mathbf{P}^1.$$

Fix $\nu = d/m > 0$ in lowest terms. Let $\mathbf{C}^\times \curvearrowright \mathfrak{g}((z))$:

$$c \cdot_\nu \gamma(z) = c^{2d\rho^\vee} \gamma(c^{2m} z) c^{-2d\rho^\vee},$$

where $2\rho^\vee = \sum_{\alpha \in \Phi^+} \alpha^\vee$.

Let $\mathfrak{g}((z))_{\nu,k}$ be the weight- $2k$ eigenspace.

In the SL_2 example, $\gamma \in \mathfrak{g}((z))_{3/2,3}$.

Lemma If γ is an eigenvector for \cdot_ν , then the induced action on $G((z))/I$ preserves $\mathcal{B}_\gamma^{\mathrm{aff}}$.

Lemma $\mathfrak{g}((z))_{\nu,0}$ is the Lie algebra of a connected reductive group $L_\nu \subseteq G((z))$. Moreover,

$$(G((z))/I)^{\mathbf{C}^\times} = \coprod_{w \in W(L_\nu) \setminus \tilde{W}} L_\nu \dot{w} I / I.$$

Fix any regular semisimple $\gamma \in \mathfrak{g}((z))_{\nu,d}$.

Springer : $\tilde{W} \curvearrowright H_c^*(\mathcal{B}_\gamma^{\mathrm{aff}}), H_{c,\mathbf{C}^\times}^*(\mathcal{B}_\gamma^{\mathrm{aff}})$.

(Sommers) If m is the Coxeter number, then:

- $L_\nu = T$ and $L_\nu \dot{w} I = \dot{w} I$.
- $(\mathcal{B}_\gamma^{\mathrm{aff}})^{\mathbf{C}^\times}$ is a finite subset of the $\dot{w} I$.
- Writing $H_{\mathbf{C}^\times}^*(\mathrm{pt}) = \mathbf{C}[\epsilon]$, we have

$$\begin{aligned} H_c^*(\mathcal{B}_\gamma^{\mathrm{aff}}) &= H_{c,\mathbf{C}^\times}^*(\mathcal{B}_\gamma^{\mathrm{aff}})|_{\epsilon \rightarrow 1} \\ &= \{\text{functions on } (\mathcal{B}_\gamma^{\mathrm{aff}})^{\mathbf{C}^\times}\}. \end{aligned}$$

- $\Pi_{\mathfrak{q}}^{(d)}(w)|_{\mathfrak{q} \rightarrow 1}$ is the W -character of $H_c^*(\mathcal{B}_\gamma^{\mathrm{aff}})$.

(Oblomkov–Yun) Filtration on $H_{c,\mathbf{C}^\times}^*|_{\epsilon \rightarrow 1}$ restores \mathfrak{q} .

(Goresky–Kottwitz–MacPherson) For general ν ,

$$(\mathcal{B}_\gamma^{\text{aff}})^{\mathbf{C}^\times} = \coprod_{w \in W(L_\nu) \setminus \tilde{W}} \text{Hess}_{\gamma, w},$$

a disjoint union of *partial Hessenberg varieties*

$$\text{Hess}_{\gamma, w} = \{gP_{\nu, w} \in L_\nu/P_{\nu, w} \mid g^{-1}\gamma g \in \dot{w}\mathfrak{J}\dot{w}^{-1}\},$$

where $P_{\nu, w} := L_\nu \cap \dot{w}I\dot{w}^{-1}$.

They are smooth.

If $\text{Hess}_{\gamma, w} \neq \emptyset$, then its codimension in $L_\nu/P_{\nu, w}$ is the number of affine roots $\alpha + k$ such that:

- $\langle \alpha, \nu\rho^\vee \rangle + k = \nu$.
- $\langle \alpha, \frac{1}{2}\rho^\vee \cdot w \rangle + k < 0$.

Conj (T) For general ν , the representation

$$W \curvearrowright H_{c, \mathbf{C}^\times}^*(\mathcal{B}_\gamma^{\text{aff}})|_{\epsilon \rightarrow 1}$$

contains a summand whose character is the $\mathfrak{q} \rightarrow 1$ limit of our earlier formula:

$$\frac{\text{sgn}(w)}{\det(1 - \mathfrak{q}w \mid \mathfrak{h})} \sum_{\chi \in \text{Irr}(W)} \mathfrak{q}^{c(\chi)\nu} D_\chi(e^{2\pi i\nu}) \chi(w).$$

Moreover, the Oblomkov–Yun filtration restores \mathfrak{q} .

Thank you for listening.