

**MATH 340: ADVANCED LINEAR ALGEBRA**  
**PROBLEM SET #7**

SPRING 2025

**Due Wednesday, April 9.** You may consult books, papers, and websites as long as you cite all sources and write up your solutions in your own words. **Updated on 4/5 at 1:30 pm, in red.**

**Problem 1.** Look up the definition of an integer partition. Let  $p(n)$  be the number of partitions of an integer  $n > 0$ . Using Jordan canonical form, show that  $p(n)$  is also the number of conjugacy classes of nilpotent  $n \times n$  matrices over  $\mathbf{C}$ .

**Problem 2.** Let

$$\mathfrak{sl}(n, F) = \{n \times n \text{ matrices over } F \text{ of trace } 0\}.$$

The notation  $\mathfrak{sl}$  stands for *special linear*.

- (1) Verify that  $\mathfrak{sl}(n, F)$  is a vector space over  $F$  of dimension  $n^2 - 1$ .
- (2) Show that an element of  $\mathfrak{sl}(2, F)$  is nilpotent if and only if **its determinant is zero**.
- (3) Using a suitable basis to identify  $\mathfrak{sl}(2, \mathbf{R})$  with  $\mathbf{R}^3$ , sketch the subset of nilpotent matrices. (No need to prove the basis is a basis.)

*In principle, the following two problems are solved in Axler's text. But it may be easier to think about them from scratch, than to start with Axler.*

**Problem 3.** Let  $V, W$  be finite-dimensional vector spaces and  $T : V \rightarrow W$  a linear map. Show that the kernel  $\ker(T^\vee)$  and the annihilator  $\text{Ann}_{W^\vee}(\text{im}(T))$  are the same subspace of  $W^\vee$ .

**Problem 4.** Keep the setup of Problem 3.

- (1) Show that

$$\dim W - \dim \ker(T^\vee) = \dim V - \dim \ker(T).$$

*Hint:* You'll need Problem 3, a dimension formula relating  $U$  and  $\text{Ann}_{W^\vee}(U)$  for some  $U \subseteq W$ , and a dimension formula relating  $\ker(T)$  and  $\text{im}(T)$ .

- (2) Deduce from (1) that

$$\dim \text{im}(T^\vee) = \dim \text{im}(T).$$

- (3) Using (2), show that the column rank and row rank of any square matrix  $M$  agree.

You may use the fact (Axler §3.132) that if  $M$  represents a linear operator  $T : V \rightarrow V$  in some basis for  $V$ , then the *transpose* matrix  $M^t$  defined by  $(M^t)_{j,i} = M_{i,j}$  represents  $T^\vee : V^\vee \rightarrow V^\vee$  in the dual basis.

**Problem 5.** Let  $V$  be a vector space over  $F$ , possibly infinite-dimensional. In each case below, show that  $T$  is a linear isomorphism without picking an explicit basis for  $V$ . You may still use the fact that a tensor product of vector spaces is spanned by pure tensors.

- (1)  $T : \{\vec{0}\} \rightarrow \{\vec{0}\} \otimes V$  defined in the only possible way.
- (2)  $T : V^{\oplus n} \rightarrow F^n \otimes V$ , where  $V^{\oplus n}$  is the  $n$ -fold direct sum of  $V$  for some  $n > 0$ , defined by

$$T(v^{(1)}, \dots, v^{(n)}) = \sum_i e_i \otimes v^{(i)},$$

where  $e_1, \dots, e_n$  is an ordered basis for  $F^n$ . *Hint:* To show injectivity, use the definition of  $e_i \otimes v^{(i)}$  as a bilinear functional and an ordered basis dual to  $(e_i)_i$ . You may use the fact that the map  $V \rightarrow (V^\vee)^\vee$  from Problem Set 6, #6 is injective for any  $V$  (even when it is not bijective).

**Problem 6.** Let  $V, W, U$  be vector spaces. The set

$$\text{Bil}(W, V \mid U) = \{\text{bilinear maps from } W \times V \text{ into } U\}$$

forms a vector space under  $(\beta + \beta')(w, v) = \beta(w, v) + \beta'(w, v)$  and  $(a \cdot \beta)(w, v) = a \cdot \beta(w, v)$ . It recovers  $\text{Bil}(W, V)$  when  $U = F$ .

For all linear  $T : W \otimes V \rightarrow U$ , let  $\beta_T : W \times V \rightarrow U$  be the bilinear map such that  $\beta_T(w, v) = T(w \otimes v)$ . Show that the map

$$B : \text{Hom}(W \otimes V, U) \rightarrow \text{Bil}(W, V \mid U) \quad \text{defined by } B(T) = \beta_T$$

is linear and injective, without picking explicit bases for the vector spaces involved. *Hint:* Again, pure tensors span  $W \otimes V$ .

*It turns out that  $B$  is an isomorphism, but starting from Axler's definition of  $W \otimes V$ , this is difficult to show without picking explicit bases for  $V$  and  $W$ .*

**Problem 7.** A bilinear form  $\beta : V \times V \rightarrow F$  is *degenerate* if and only if there is some nonzero  $v \in V$  such that either  $\beta(v, -)$  or  $\beta(-, v)$  is the zero functional on  $V$ . It is *nondegenerate* otherwise. Now set  $V = F[x]$ . Show that:

- (1) If  $\beta(p, q) = \int_0^1 p(x)q(x) dx$ , then  $\beta$  is nondegenerate.
- (2) If  $\beta(p, q) = p(1)q(1)$ , then  $\beta$  is degenerate.

**Problem 8.** Show that for all  $n \geq 2$ , there is a bilinear form  $\beta$  on  $F^n$  such that

$$\beta(w, v) \neq 0 \text{ for some } w, v \in F^n, \quad \text{but } \beta(v, v) = 0 \text{ for all } v.$$

*Hint:* Take  $\beta(w, v) = w^t M v$  for some carefully chosen  $n \times n$  matrix  $M$ .